arXiv:1202.4909v2 [math.NT] 6 Mar 2012

REPRESENTATION ON AVERAGE OF BINARY QUADRATIC FORMS BY AN INTEGRAL QUATERNARY QUADRATIC FORM

RAINER SCHULZE-PILLOT

Abstract. We investigate the average of the number of representations of an integral binary quadratic form of fixed discriminant $-d$ by certain integral quaternary quadratic forms. We also find an estimate on the number of binary forms of fixed discriminant $-d$ which are represented by the given quaternary form.

1. Introduction.

If A is an integral positive definite symmetric $(m \times m)$ matrix one calls a solution $X \in M(m \times n, \mathbb{Z})$ of the system of quadratic equations of the form

$$Q_A(X) = {}^tXAX = T$$

where T is another (half-) integral symmetric matrix of size $n \leq m$ an (integral) representation of T by S. It is well known that the local global principle of Minkowski and Hasse is not valid for integral representations, but if m is large enough compared to n one can prove that a positive definite T which is represented by S over all $\mathbb{Z}_p$ and is large enough in a suitable sense is indeed represented by S over the rational integers $\mathbb{Z}$, at least under some mild additional conditions. The bound on the size of m necessary for this has recently been pushed down to $m \geq n+3$, again under suitable additional conditions, in [5], see also [10] for an attempt to optimize those additional conditions. The case $m = n+2$ brings some limitations due to the existence of the so called spinor exceptions (see [7, 6]), taking these into account a result of the desired type could be reached in [4] for $n = 1, m = 3$, i. e. for representations of sufficiently large numbers by ternary forms. The case of $m = n+2$ appeared to be rather intractable for $n \geq 2$ until recently, when Einsiedler, Lindenstrauss, Michel, and Venkatesh obtained a result for a certain subset of T by adapting Linnik's ergodic method, which was invented to deal with the ternary case, to the problem of representation of binary forms by quaternary forms. They also obtained a result on representation of T "on average" under the assumption of the so called Linnik condition.

A similar, but somewhat different average result follows, at least for S related to orders of square free level in quaternion algebras, almost immediately from a result of Böcherer and the author from [2], without assuming the Linnik condition mentioned, however, at the time when [2] was published there seemed to be little interest in such average results. The purpose of this note is to take the occasion to show how this can be done. In this first version we restrict to the above mentioned class of S. It is likely that the argument can be extended to other S, if so, this will be done in a second version which then will replace this version. In the final section we discuss some speculations about estimates for averages of Fourier coefficients of Siegel modular forms that are related to the results obtained here.

I heard of the results of Einsiedler, Lindenstrauss, Michel, and Venkatesh in the talk of Michel at the International Colloquium on Automorphic Representations and L-functions 2012 at the Tata Institute of Fundamental Research, Mumbai, where this note was also written during a stay following that colloquium. I thank the institute, in particular Dipendra Prasad, for its resp. his hospitality.

2. Lattices in quaternion algebras

Let D be a quaternion algebra over $\mathbb{Q}$ which is ramified at infinity and at the primes dividing the square free integer N_1 (which then has an odd number of prime factors) and isomorphic to the ring of 2×2 matrices at the other primes. We denote by n the reduced norm and by tr the reduced trace and view D as a quadratic space with quadratic form n and associated symmetric bilinear form $b(x, y) = \mathrm{tr}(\mathrm{x\bar{y}})$, where $y \mapsto \bar{y}$ is the usual involution (quaternionic conjugation) on D. Let R be an Eichler order of square free level $N = N_1 N_2$, i.e., $R_p := R \otimes \mathbb{Z}_p$ is isomorphic (in fact conjugate) to the ring of matrices of the form $\left(\begin{smallmatrix} * & * \\ p* & * \end{smallmatrix}\right)$ where $*$ indicates an element of $\mathbb{Z}_p$ for all primes dividing N_2, and R_p is a maximal order in $D_p := D \otimes \mathbb{Q}_p$ at the other primes. It is well known that the adelization $D_{\mathbb{A}}^{\times}$ has a finite double coset decomposition $D_{\mathbb{A}}^{\times} = \cup_{j=1}^{h} D_{\mathbb{Q}}^{\times} y_j R_{\mathbb{A}}^{\times}$, where the adeles y_j have trivial infinity component and are of norm 1 at all places. The R-ideals $I_j := R(y_j)^{-1}$ are then a set of representatives of the classes of left R ideals modulo principal ideals (on the right) whose left order is equal to R, and more generally the $I_{ij} = y_i R(y_j)^{-1}$ represent the classes of ideals with left order $R_i := y_i R(y_i)^{-1}$ and right order R_j. It is well known that if we let N_1, N_2 vary over all decompositions $N = N_1 N_2$ of N where N_1 has an odd number of prime factors, the quadratic lattices (I_{ij}, n) run through all isometry classes of integral quaternary quadratic lattices of discriminant N^2 and level N, with possible multiple appearances of isometry classes. For any fixed decomposition all the lattices arising in this way form a genus of quadratic lattices consisting of a single spinor genus.

We further denote for $s \mid N$ by $I_{ij}^{\#,s}$ the lattice obtained as $I_{ij}^{\#} \cap \mathbb{Z}[\frac{1}{s}]$, where # indicates taking the dual lattice, and by $I_{ij}^{*,s}$ the same lattice with the quadratic forms scaled by the factor s. We notice that this last lattice in the same genus as the I_{ij} and is hence isometric to some $I_{i'j'}$. We call the $I_{ij}^{*,s}$ the rescaled partial duals of I_{ij} and write $\mathrm{rpd}(\mathrm{I_{ij}}, \mathrm{N'})$ for the set of all $I_{ij}^{*,s}$ with $s \mid N'$, where N' is some fixed divisor of N. For a positive definite matrix $T = (t_{kl})_{k,l=1..2} \in M^{\mathrm{sym}}(2, \mathbb{Z})$ we write

$$r(I_{ij}, T) := |\{(x, y) \in I_{ij} \mid \begin{pmatrix} n(x) & b(x,y) \\ b(x,y) & n(y) \end{pmatrix} = T\}|$$

for the number of representations of T by the quadratic lattice (I_{ij}, n) and $r^*(I_{ij}, T)$ for the number of primitive representations, i.e., representations (x, y) where $\mathbb{Z}x + \mathbb{Z}y$ is a direct (but not necessarily orthogonal) summand of I_{ij} as a $\mathbb{Z}$-module.
We write

$$L_i := \{x \in \mathbb{Z} \cdot 1 + 2R_i \mid \mathrm{tr(x)} = 0\}$$

and $r(L_i, t) = |\{x \in L_i \mid n(x_i) = t\}|$ for a positive integer t. The quadratic lattice (L_i, n) has level $4N$ and discriminant $32N^2$. It is easily checked that for a fixed decomposition $N = N_1 N_2$ all the L_i are in the same genus of quadratic lattices and that this genus contains only one spinor genus.

3. Averages of representation numbers

With notations as above we write

$$r(I_{ij}, d) = \sum_T \frac{r(I_{ij}, T)}{\epsilon(T)},$$

where the sum is over a set of $SL_2(\mathbb{Z})$-equivalence classes of integral symmetric matrices $T = \left(\begin{smallmatrix} 2a & b \\ b & 2c \end{smallmatrix}\right)$ of (signed) discriminant $b^2 - 4ac = -d$ and where $\epsilon(T)$ denotes the number of proper automorphisms (or units) of T, i. e. the number of $U \in SL_2(\mathbb{Z})$ with ${}^tUTU = T$. The primitive average $r^*(I_{ij}, d)$ is defined analogously by using the primitive representation numbers $r^*(I_{ij}, T)$.
We want to use Theorem 2.1 from [2] which gives:

If $-d$ is a discriminant and $N_d := \frac{N}{\gcd(N,d)}$ one has

$$\sum_{s \mid N_d} \sum_{m^2 \mid d} r^*(I_{ij}^{*,s}, \frac{d}{m^2}) = r(L_i, d) r(L_j, d). \tag{3.1}$$

If N is prime this simplifies to

$$\sum_{m^2 \mid d} r^*(I_{ij}, \frac{d}{m^2}) = \frac{1}{2} r(L_i, d) r(L_j, d). \tag{3.2}$$

In particular, if in addition $-d$ is fundamental the sum over $m^2 \mid d$ can be omitted in both formulas.

As an easy consequence we obtain:

Theorem 1. *Let* $\epsilon > 0$. *Then there are constants* $C_i = C_i(\epsilon, N)$ *for* $i = 1, 2$ *such that:*

a) *If N is prime one has $r(I_{ij},d) \geq C_2 d^{1-\epsilon}$ for all $d > C_1$ which are represented primitively over all $\mathbb{Z}_p$ by L_1 (and hence by all the L_i).*
b) *If N is composite and $N' \mid N$ a fixed divisor of N, one has $r(\mathrm{rpd}(\mathrm{I_{ij}},\mathrm{N'}),\mathrm{d}) \geq \mathrm{C_2 d^{1-\epsilon}}$ for all $d > C_1$ with $N_d = N'$ which are represented primitively over all $\mathbb{Z}_p$ by L_1 (and hence by all the L_i). Here we write $r(\mathrm{rpd}(\mathrm{I_{ij}},\mathrm{N'}),\mathrm{d}) = \sum_{\mathrm{s}\mid\mathrm{N'}} \mathrm{r}(\mathrm{I_{ij}^{*,s}},\mathrm{d})$.*

If one restricts here to d for which $-d$ is fundamental the local condition is satisfied if and only if p is not split in $\mathbb{Q}(\sqrt{-d})$ for all $p \mid N_1$ and is not inert in this quadratic extension for all $p \mid N_2$.

Proof. The easiest case is $-d$ fundamental, in which case the assertion is a direct consequence of the formula (3.2) if N is prime resp. (3.1) if N is composite from [2] quoted above and the result of [4] giving an asymptotic formula for the representation numbers of ternary quadratic forms (the possible obstruction by spinor exceptions plays no role here since the L_i are in a genus consisting of one spinor genus). If $-d$ is not fundamental we use the asymptotic formula for primitive representations of ternary lattices (by assumption d is locally everywhere primitively representable by L_i, L_j), which gives the same rate of growth, and recall that the proof of Theorem 2.1 of [2] associates to a pair (up to sign) of primitive vectors of length d in L_i resp. L_j a primitive representation of a binary matrix T counted in the left hand side of equation (3.2) in the case that N is prime, and a pair of primitive representations of a binary matrix T by two (different) lattices $I_{ij}^{*,s}$ counted in the left hand side of equation(3.1), so that the term $r^*(\mathrm{rpd}(\mathrm{I_{ij}},\mathrm{N'}),\mathrm{d}) \leq \mathrm{r}(\mathrm{rpd}(\mathrm{I_{ij}},\mathrm{N'}),\mathrm{d})$ in the sum for $r(L_i,d)r(L_j,d)$ above satisfies the required estimate. $\square$

Remark. *The appearance of the set of all rescaled partial duals $I_{ij}^{*,s}$ with $s \mid N'$ of I_{ij} for the d not divisible by N (and with $N_d = N'$) in the case of composite N is somewhat unsatisfactory. The number of lattices involved is obviously $2^{\omega(N')}$, but the set of distinct isometry classes of these lattices may be smaller. The argument from* [2] *would allow to replace representations of T by $I_{ij}^{*,s}$ by representations of sT by I_{ij} itself, if one likes that better.*

4. AVERAGE ASYMPTOTIC FORMULA AND APPLICATIONS

We restrict attention now to fundamental discriminants $-d$.

Proposition 2. *Let $-d$ be a fundamental discriminant and $N_d = \frac{N}{\gcd(N,d)}$. Then we have:*

$$h(-d)\sigma_0(N_d)r(\mathrm{gen(R)},\mathrm{T}) = (\mathrm{r}(\mathrm{gen}(\mathrm{L_i}),\mathrm{d})^2, \tag{4.1}$$

where σ_0 denotes the number of divisors function.

Proof. It is easily seen that Siegel's weighted average $r(\mathrm{gen(R),T})$ of the numbers of representations of T by the classes of quadratic lattices in the genus of (R, n) (i.e., in the genus consisting of the I_{ij}) is independent of T, in particular it has the same value for T in all genera of binary quadratic forms of discriminant $-d$. Upon summation over the i, j, s with weights $\frac{1}{e_i e_j}$, where e_i is the number of units of the order R_i, the assertion is then seen to be a direct consequence of equation (3.1). Of course, the assertion could also be established by a direct computation of local densities. □

From [4] we have

$$r(L_i, d)r(L_j, d) = (r(gen(L_1, d)))^2 + O(d^{1+\epsilon-\frac{1}{16}}), \tag{4.2}$$

where we used the best currently known estimate for the error term in the asymptotic formula for $r(L_i, d)$, see [1]. Putting this together with the proposition and equation (3.1) we obtain

Theorem 3 (Average asymptotic formula)**.** *Let $N' \mid N$ be fixed and let $-d$ run over fundamental discriminants with $N_d = N'$. Then we have for all $\epsilon > 0$*

$$\begin{aligned} r(\mathrm{rpd(I_{ij}, N'), d}) &= (r(\mathrm{gen(L_1), d}))^2 + \mathrm{O(d^{1+\epsilon-\frac{1}{16}})} \\ &= h(-d)\sigma_0(N_d)r(\mathrm{gen(R), T}) + \mathrm{O(d^{1+\epsilon-\frac{1}{16}})}, \qquad (4.3) \end{aligned}$$

where T is the matrix of any integral binary quadratic form of discriminant $-d$.
If N is prime we have the better result (independent of $\gcd(N, d)$*)*

$$r(I_{ij}, d) = h(-d)r(\mathrm{gen(R), T}) + \mathrm{O(d^{1+\epsilon-\frac{1}{16}})}. \tag{4.4}$$

In order to obtain from this a result about representations of individual forms T we denote by $\mu(R)$ the mass of the genus of the quadratic lattice R and by $o_{\max}(R)$ the maximal order of the group of automorphisms of a lattice in the genus of R. With this notation we have:

Theorem 4. *Let N' be a fixed divisor of N and let $-d$ run through fundamental discriminants with $N_d = N'$ such that d is represented over all $\mathbb{Z}_p$ by the lattices in the genus of the L_i (equivalently, binary quadratic forms of discriminant $-d$ are represented over all $\mathbb{Z}_p$ by the I_{ij}). Then for all $\kappa < 1$ there is a constant C_3 such that for all $d \geq C_3$ as above the number ν of binary quadratic forms T of discriminant $-d$ with* $r(\mathrm{rpd(I_{ij}, N'), T}) \neq 0$ *satisfies*

$$\nu \geq \kappa \frac{\sigma_0(N_d)h(-d)}{\mu(R)o_{\max}(R)}.$$

In particular, for d large enough a positive proportion of the classes of binary quadratic forms of discriminant $-d$ is represented by at least one among the rescaled partial dual lattices of I_{ij} comprising the set $\mathrm{rpd(I_{ij}, N_d)}$*.*

If N is prime we let ν' denote the number of T as above represented by I_{ij} itself, get

$$\nu' \geq \kappa \frac{h(-d)}{\mu(R) o_{\max}(R)}$$

for sufficiently large d and conclude that for d large enough a positive proportion of the classes of binary quadratic forms of discriminant $-d$ is represented by I_{ij} itself.

Proof. From Siegel's mass formula we see that we must have

$$r(\mathrm{rpd}(\mathrm{I_{ij}}, \mathrm{N_d}), \mathrm{T}) \leq \mathrm{o_{max}(R)}\mu(\mathrm{R})\mathrm{r}(\mathrm{gen}(\mathrm{R}), \mathrm{T}), \tag{4.5}$$

since all terms in the weighted sum for $r(\mathrm{gen}(\mathrm{R}), \mathrm{T})$ are non negative. Summing over the T of discriminant $-d$ and taking into account the asymptotic formulas from Theorem 3 we obtain the assertion (we use the asymptotic formula to conclude that the sum of the main term and the error term is for $\kappa < 1$ and sufficiently large d at least as large as κ times the main term). ☐

Remark. a) *The "sufficiently large" in the theorem is not effective since our use of the asymptotic formula relies on the fact that $h(-d)$ is at least of the order of $d^{\frac{1}{2}-\epsilon}$ for all $\epsilon > 0$, which estimate is well known to be ineffective.*

b) *Our method appears not to be suitable to guarantee the existence of T which is represented by I_{ij} and lies in a given small (say of size d^{δ}) subset of the class group.*

c) *If the number of represented forms T is as small as it can be the representation number $r(I_{ij}, T)$ (or $r(\mathrm{rpd}(\mathrm{I_{ij}}, \mathrm{N'}), \mathrm{T})$ has to be rather large, i. e. of the order of magnitude of the number $r(\mathrm{gen}(\mathrm{R}), \mathrm{T})$ from Siegel's mass formula (in fact, even larger).*

d) *In contrast to most other results about representation of quadratic forms of rank > 1 by quadratic forms our result does not at all involve the minimum of the binary quadratic form T to be represented, at least not explicitly.*

5. SPECULATIONS

The results from [2] quoted above where used there to compute averages of Fourier coefficients of special Siegel modular forms of degree 2, the Yoshida liftings. These averages appear there as the coefficients in the Dirichlet series of Koecher and Maaß associated to such a Siegel modular form. An immediate consequence is the following:

Proposition 5. *Let $F \in M_k^{(2)}(\Gamma_0(N))$ be the Yoshida lifting of a pair f_1, f_2 of primitive cuspidal elliptic Hecke eigenforms of weights $k_1 = 2, k_2 = 2k - 2$ and level N and denote by*

$$a(F, d) = \sum_T \frac{a(F, T)}{\epsilon(T)}$$

the $d-th$ coefficient of its Koecher-Maaß series, where $a(F,T)$ is the T-th Fourier coefficient of F, the sum is over a set of $SL_2(\mathbb{Z})$-equivalence classes of integral symmetric matrices $T = \left(\begin{smallmatrix} 2a & b \\ b & 2c \end{smallmatrix}\right)$ of (signed) discriminant $b^2 - 4ac = -d$, and where $\epsilon(T)$ denotes the number of proper automorphisms (or units) of T.
Then for any $\epsilon > 0$ there is a constant C such that one has for all d

$$a(F,d) \leq C d^{\frac{k}{2}-\frac{1}{8}+\epsilon}. \tag{5.1}$$

If above we replace the cusp form f_1 by the (appropriate) Eisenstein series of weight 2 and level N we obtain

$$a(F,d) \leq C d^{\frac{k}{2}-\frac{1}{16}+\epsilon}. \tag{5.2}$$

Remark.

- *If one assumes the generalized Ramanujan conjecture for cusp forms of half integral weight in the complement of the one-dimensional theta series the exponents above would improve to $\frac{k}{2} - \frac{1}{2} + \epsilon$ in the first case and to $\frac{k}{2} - \frac{1}{4} + \epsilon$ in the second case above.*
- *In the second case above the form F is in the space of Maaß or Saito-Kurokawa lifts, so that its Fourier coefficient at T depends only on the determinant of T and the summation over T just multiplies the individual coefficient by the number of classes of T of determinant d.*
- *Similar estimates can be obtained for vector valued Yoshida liftings associated to a pair of elliptic cusp forms k_1, k_2 with $k_1 > 2, k_2 > 2$, using the results of* [3]. *If such a vector valued lifting has transformation type $\mathrm{Sym}^j \otimes \det^k$ we obtain an exponent $\frac{j+k}{2} - 2\delta + \epsilon$ at d with $\delta = \frac{1}{16}$ unconditionally and $\delta = \frac{1}{4}$ under the assumption of the generalized Ramanujan conjecture for cusp forms of half integral weight $\geq \frac{5}{2}$*
- *A well known conjecture of Böcherer relates $a(F,d)$ to the square of the central critical value of the twist by χ_d of the spin L-function of F. If we assume this conjecture and a subconvexity bound for this L-value (with respect to the conductor d of the character) we get bounds for $a(F,d)$ of the above type; if we assume Böcherer's conjecture and the Lindelöf hypothesis (again with respect to d) for this L-value we get the version given in a) which assumes the generalized Ramanujan conjecture for cusp forms of half integral weight. In the special case of the Yoshida liftings both Böcherer's conjecture and the required subconvexity estimate are known.*
- *In* [8] *the Fourier coefficient at T with $\det(T) = d$ of a cusp form of weight k and degree 2 for the full Siegel modular group is estimated to be $O(d^{\frac{k}{2}-\frac{13}{36}+\epsilon}$. If we multiply this estimate by the number of integral equivalence classes of T of determinant d we obtain an exponent of $\frac{k}{2} + \frac{5}{36} + \epsilon$. The conjecture of Resnikoff and*

Saldaña [9] would give an exponent of $\frac{k}{2} - \frac{3}{4} + \epsilon$ *for the individual Fourier coefficient and hence an exponent of* $\frac{k}{2} - \frac{1}{4} + \epsilon$ *for the sum* $a(F, d)$*, which bound should be compared with the conjectural bound* $\frac{k}{2} - \frac{1}{2} + \epsilon$ *obtained above for Yoshida liftings outside the Maaß space under the assumption of the generalized Ramanujan conjecture for (good) cusp forms of half integral weight, or for general cusp forms orthogonal to the Maaß space under the assumption of Böcherer's conjecture and the Lindelöf conjecture for the twisted spin L-function. If we consider only forms in the complement of the Maaß space we see hence that our result for Yoshida liftings gives results for the averaged coefficient* $a(F, d)$ *that are significantly better than those obtained by multiplying the estimate for the individual Fourier coefficient by the number of summands.*

We might summarize the statements of the remark above in the speculation that an estimate of the type

$$a(F, d) \leq Cd^{\frac{k}{2} - \delta + \epsilon} \tag{5.3}$$

with some $\delta > 0$ could be true and perhaps provable for a general Siegel modular cusp form of degree 2, including as in part c) of the above remark the vector valued case. We might also speculate that similar estimates may hold for averaged Fourier coefficients of higher degree Siegel cusp forms. This problem will be the subject of future work.

References

[1] V. Blomer, G. Harcos: Hybrid bounds for twisted L-functions. J. Reine Angew. Math. **621** (2008), 53-79

[2] S. Böcherer, R. Schulze-Pillot, The Dirichlet series of Koecher and Maaß and modular forms of weight $\frac{3}{2}$. *Math. Z.* **209** (1992), no. 2, 273–287.

[3] S. Böcherer, R. Schulze-Pillot, Mellin transforms of vector valued theta series attached to quaternion algebras, *Math. Nachr.* **169** (1994), 31–57.

[4] W. Duke, R. Schulze-Pillot: Representation of integers by positive ternary quadratic forms and equidistribution of lattice points on ellipsoids. *Invent. Math.* **99** (1990), 49–57.

[5] J. Ellenberg, A. Venkatesh: Local-global principles for representations of quadratic forms. *Invent. Math.* **171** (2008), 257–279.

[6] J. S. Hsia: Representations by spinor genera. *Pacific J. Math.* **63** (1976), 147–152.

[7] M. Kneser: Darstellungsmaße indefiniter quadratischer Formen. *Math. Z.* **77** (1961), 188–194.

[8] W. Kohnen: Estimates for Fourier coefficients of Siegel cusp forms of degree two, II, *Nagoya Math. J.* **128** (1992), 71–76

[9] H. L. Resnikoff and R. L. Saldaña: Some properties of Fourier coefficients of Eisenstein series of degree two, *J. reine angew. Math.* **265** (1974) 90–109.

[10] R. Schulze-Pillot: Local conditions for global representations of quadratic forms, *Acta Arith.* **138** (2009), 289–299

Rainer Schulze-Pillot
Fachrichtung 6.1 Mathematik, Universität des Saarlandes (Geb. E2.4)
Postfach 151150, 66041 Saarbrücken, Germany
email: schulzep@math.uni-sb.de